Mouse Anti-HCV Core Protein Monoclonal Antibody Datasheet

Product Name: mAb anti-HCV Core Protein Clone No.: C3

Catalogue No.: MO-I40015E Quantity: 0.5 mg/vial

Description: Mouse monoclonal antibody (mAb) to

Human hepatitis C Virus (HCV) core protein

Purification: Protein G affinity purified

Product Type: Primary antibody

Target Protein: Human hepatitis C virus (HCV) core protein

Immunogen: Recombinant Chimeric Hepatitis C virus

(HCV) polyprotein (555 amino acids)

Fusion Sp2/0-Ag14

Myeloma:

Specificity: This mAb is reactive to recombinant core

protein C + envelope protein M (residues 1-142 on HCV polyprotein) and synthetic core

protein C (residues 1-61 on HCV

polyprotein), and recombinant chimeric

HCV polyprotein (60kDa).

Species Human hepatitis C virus, others not tested

Reactivity:

Cross - No cross reaction with recombinant NS-3

Reactivity: protein (1252nd -1477th amino acid),

synthetic NS-3 protein (1378th -1458th amino acid) and synthetic NS-4a protein

(1689th -1735th amino acid).

Host / Isotype: Mouse, IgG1 Kappa

Formulation: Lyophilized from a solution in 0.01M PBS,

pH 7.0

Reconstitution: Double distillated water is recommended

to adjust the final concentration to

1.00mg/mL.

Storage: Store at -20°C

Research Area: Virology

Background: Hepatitis C virus (HCV) causes chronic

hepatitis and liver cirrhosis in human through blood and body fluid transmission.

HCV has a positive sense single RNA

genome enclosed in the nucleocapsid made of Core Protein (Capsid Protein). The nucleocapsid is covered by an envelope made of lipoproteins (E1 and E2). The 9.6 kb HCV genome has a single open-reading frame, which is to be translated into a

single polyprotein. HCV viral proteins are produced after processing the polyprotein. Genes for core protein and envelop

proteins are located adjacently at the 5'end of HCV genome, followed by genes for non-structural proteins including NS2, NS3,

NS4A, NS4B, NS5, NS5A and NS5B.

Applications: ELISA: The mAb is reactive to HCV core

protein and not reactive with HCV non-

structural protein.

Western Blot: The antibody when used at

concentration of $0.1\mu g/ml$, will allow visualization of $0.1\mu g/lane$ recombinant core protein C + envelope protein M (residues 1-142 on HCV polyprotein), $0.5\mu g/lane$ synthetic core protein C (residues 1-61 on HCV polyprotein), and $0.1\mu g/lane$ recombinant 60kDa chimeric HCV polyprotein. The mAb works on blots transferred from both reducing and non-reducing PAGE gel. The mAb has been

vitro translated HCV core protein.

References: If research is published using this product,

please inform Anogen in order to cite the reference on this datasheet. Anogen will provide one unit of product in the same

used successfully in the recognition of in-

category as gratitude.

This product is for LABORATORY RESEARCH USE and further manufacture ONLY, and cannot be administrated to human and animals for use in diagnostic and therapeutic procedures.

Manufactured by ANOGEN - A Division of YES Biotech Laboratories Ltd.

Page 1 of 1 page(s)

To place order, please contact us by phone, fax or by email: info@anogen.ca,
or with our secure online store: www.anogen.net